

REAL-WORLD TREATMENT EXPERIENCE OF SINGLE TABLET DOLUTEGRAVIR/LAMIVUDINE IN THE US: RESULTS FROM THE TANDEM STUDY

S Schneider,¹ C Burke,² D Ward³, P Benson⁴, C Donovan⁵, G Harper⁶, D Merrill⁵, A Metzner⁵, K Mycock⁶, J Patarroyo⁵, A Oglesby⁵

¹Long Beach Education and Research Consultants, Long Beach, CA, USA; ²CAN Community Health, St Petersburg, FL, USA; ³Dupont Circle Physicians Group, Washington DC, USA; ⁴Be Well Medical Center, Berkley, MI, USA; ⁵ViiV Healthcare, Research Triangle Park, NC, USA; ⁶Adelphi Real World, Bollington, Cheshire, UK

Key Takeaways

- TANDEM aimed to characterize real-world prescribing behaviors and treatment outcomes in the United States (US).
- Out of a total population of 469, 318 received dolutegravir/ lamivudine (DTG/3TC), of whom 126 were naïve to antiretroviral therapy (ART), and 192 were stable-switch (SS).
- After an average follow-up time of >1 year, 93.7% of treatment-naïve PLWH achieved virological suppression while receiving DTG/3TC and 95.8% of SS PLWH maintained virological suppression on DTG/3TC following switch.
- TANDEM reflects results from phase 3 clinical trials demonstrating DTG/3TC is an effective and well tolerated regimen with few treatment discontinuations.
- Primary reasons for initiation of DTG/3TC were to avoid long term toxicities (both cohorts), simplification/ streamlining of treatment (for SS sub-cohort) and convenience (for treatment-naïve sub-cohort).

Introduction

- Treatment for People Living With HIV-1 (PLWH) continues to advance with a two-drug regimen (2DR) approach. ^[1]
- DTG/3TC is indicated as a 2DR for both treatment-naïve and virally suppressed PLWH. ^[2]
- Treatment with 2DRs has the potential to decrease lifetime cumulative drug exposure and long-term toxicities associated with multiple antiretrovirals. ^[3]
- DTG/3TC demonstrated non-inferiority vs. DTG+ tenofovir disoproxil fumarate/emtricitabine out to 3 years in the phase 3, double-blinded GEMINI-1 and GEMINI-2 trials.^[4]
- Despite sustained virologic efficacy for DTG-based 2DRs observed in clinical trials, there is limited evidence in US-based real-world clinical settings. The TANDEM study aimed to characterize real-world prescribing behaviors and treatment outcomes in the US.

Methods

- TANDEM was a US-based, retrospective chart review. 24 sites abstracted data from medical charts of PLWH who were initiated on DTG/3TC or dolutegravir / rilpivirine (DTG/RPV) prior to Sept/30/2020, with a minimum clinical follow-up of six months.
 - Minimum follow-up could include time post-discontinuation of either regimen.
- Clinical characteristics, treatment history and post-initiation outcomes were abstracted.
- Analyses were descriptive and no formal hypotheses were tested.
- Missing data were not imputed. Descriptive analyses were performed in IBM® SPSS® Data Collection Survey Reporter v7.5 software.
- Time to event outcomes were calculated using Kaplan-Meier estimators conducted in StataCorp, 2015. Stata statistical software: Release 16 (College Station, TX, StataCorp LP).

Inclusion Criteria

- ≥18 years old;
- Have a diagnosis of HIV-1 infection;
- Have a history of ART therapy consisting of 2DR;
- One of the antiretroviral therapies must be either DTG/3TC or DTG/RPV as a single-tablet regimen (STR);
- DTG/3TC cohort:
 - Must have been initiated after 1st May 2019
 - Upon initiation, PLWH must have been either naïve (N) to ART or virologically suppressed defined as having HIV-1 RNA <50 copies/mL, on a stable ART regimen for ≥3 months upon DTG-based 2DR initiation (SS).
- Results for the DTG/RPV cohort will be reported separately
- At least 6 months of clinical follow-up after initiation of DTG-based 2DR.

Results

PLWH Characteristics

- From an overall sample of 469 PLWH, 151 received DTG/RPV and 318 received DTG/3TC, of whom 126 were treatment-naïve and 192 were SS. 48.4% of treatment-naïve PLWH received DTG/3TC as part of a test and treat paradigm.
- Demographics of the treatment-naïve and SS DTG/3TC cohort are described in **Table 1**.
 - The treatment-naïve cohort had a younger mean age compared to the SS cohort (37.4 vs 49.1 years).
- Common treatment considerations for the SS cohort were comorbidities (25.0%) and polypharmacy (12.5%), and for the treatment-naïve cohort were limited access to healthcare (12.7%) and comorbidities (9.5%) (**Table 2**).
 - At the data cut-off, median time on DTG/3TC was 1.3 years (treatment-naïve) [IQR 0.8,1.8] and 1.6 years (SS) [IQR 1.2, 1.8].
 - In the SS cohort, PLWH had received a median of 8.6 years prior ART and 66.1% received ≥2 prior regimens.
- From the HCP perspective, the most common reasons for initiating DTG/3TC were avoidance of long-term toxicities in both treatment-naïve (32.5%) and SS (27.1%) cohorts (**Figure 1**).
 - In the SS cohort, simplification/ streamlining (25.0%) and managing existing toxicities/ intolerance issues (12.5%) were also of high importance.
 - In the treatment-naïve cohort, convenience (15.9%), PLWH preference (12.7%) and weight gain (7.9%) were also common considerations.


Table 1. DTG/3TC PLWH Demographics

	DTG/3TC Naïve (N) n=126	DTG/3TC Stable Switch (SS) n=192
Age (years)		
Mean (SD)	37.4 (±12.7)	49.1 (±12.4)
Assigned Sex at Birth, n (%)		
Male	111 (88.1%)	158 (82.3%)
Current gender identity, n (%)		
Cis-male	103 (81.7%)	148 (77.1%)
Cis-female	15 (11.9%)	34 (17.7%)
Trans-female	4 (3.2%)	2 (1.0%)
Race, n (%)		
White/ Caucasian	77 (61.1%)	124 (64.6%)
Black	36 (28.6%)	57 (29.7%)
Pacific islander	3 (2.4%)	0 (0.0%)
Asian	2 (1.6%)	2 (1.0%)
Not specified	5 (4.0%)	9 (4.7%)
Ethnicity, n (%)		
Hispanic / Latinx, n (%)	50 (39.7%)	48 (25.0%)
Current Insurance Coverage, n (%)		
Employee provided/ sponsored insurance	34 (27.0%)	64 (33.3%)
Privately arrange insurance	23 (18.3%)	41 (21.4%)
Medicare	8 (6.3%)	30 (15.6%)
Medicaid	22 (17.5%)	26 (13.5%)
Health insurance exchange plan	16 (12.7%)	14 (7.3%)
AIDS Drug Assistance Program (ADAP)	20 (15.9%)	14 (7.3%)
No insurance coverage	3 (2.4%)	3 (1.6%)

Table 2. DTG/3TC PLWH Characteristics

	DTG/3TC Naïve (N) n=126	DTG/3TC Stable Switch (SS) n=192
Median time on DTG/3TC, years (IQR)	1.3 (0.8,1.8)	1.6 (1.2,1.8)
Median prior ART duration, years (IQR)	-	8.6 (3.5,15.3)
Number of prior ART regimens, n (%)		
1	-	65 (33.9%)
≥2	-	127 (66.1%)
Relevant Treatment Considerations, n (%)		
Comorbidities	12 (9.5%)	48 (25.0%)
Polypharmacy	2 (1.6%)	24 (12.5%)
Health insurance issues or changes	8 (6.3%)	19 (9.9%)
Mental Health issues	9 (7.1%)	16 (8.3%)
Limited access to healthcare	16 (12.7%)	10 (5.2%)
Substance abuse	6 (4.8%)	7 (3.6%)
Affordability of HIV medication	5 (4.0%)	11 (5.7%)
Adherence issues	4 (3.2%)	6 (3.1%)
Job Instability	7 (5.6%)	6 (3.1%)
Difficult work and/ or family schedule	5 (4.0%)	4 (2.1%)
Low Health Literacy	7 (5.6%)	4 (2.1%)
Homelessness/ Unstable living conditions	2 (1.6%)	3 (1.6%)
Food insecurity	2 (1.6%)	1 (0.5%)
Other	3 (2.4%)	3 (1.6%)
None	56 (44.4%)	98 (51.0%)

Figure 1. HCP Reasons for Initiating PLWH on DTG/3TC



Virological Outcomes

- 95.8% of SS PLWH maintained suppression while receiving DTG/3TC. Six of the 8 SS PLWH who then became detectable remained on DTG/3TC during the follow-up period, with 2 lost to follow-up (**Table 3**).
- 93.7% of treatment-naïve PLWH achieved virological suppression and 83.3% remained suppressed.
- Five of the 6 treatment-naïve PLWH who rebounded remained on DTG/3TC, with 1 lost to follow-up. At point of abstraction, these 5 treatment-naïve PLWH had been on DTG/3TC for a median duration of 3.6 months post-rebound (IQR: 3.3 – 12.9 months) and 11.6 months total.
- N=1 treatment-naïve and N=3 SS PLWH discontinued DTG/3TC by the data cut-off.

Table 3. Virological Outcomes

	DTG/3TC Naïve (N) n=126	DTG/3TC Stable Switch (SS) n=192
Virological status of SS PLWH, n (%)		
Remained virologically suppressed	-	184 (95.8%)
Became detectable, remained on DTG/3TC and resuppressed	-	4 (2.1%)
Became detectable, remained on DTG/3TC and not resuppressed	-	2 (1.0%)
Don't know	-	2 (1.0%)
Virological status of N PLWH, n (%)		
Became virologically suppressed	118 (93.7%)	-
- Then rebounded	6 (4.8%)	-
- Then remained suppressed	105 (83.3%)	-
- Then lost to follow-up	7 (5.6%)	-
Remained virologically detectable	3 (2.4%)	-
Don't know/ lost to follow-up	5 (4.0%)	-
Time from DTG/3TC initiation to PLWH becoming suppressed (weeks)		
Median (IQR)	10.4 (5.7,19.1)	-
Discontinuation, n (%)		
Discontinued DTG/3TC	1 (0.8%)	3 (1.6%)
Ongoing DTG/3TC	123 (97.6%)	188 (97.9%)
Unknown/ lost to follow-up	2 (1.6%)	1 (0.5%)
Time from initiation to discontinuation (weeks)*		
Median (IQR)	60.9 (60.9, 60.9)	29.6 (21.5, 74.3)
Reason for discontinuation, n (%)*		
Viremia (persistent low level/ viral blips)	1 (0.8%)	0 (0.0%)
Toxicity/ intolerance	0 (0.0%)	1 (0.5%)
Concerns about weight gain	0 (0.0%)	1 (0.5%)
PLWH preference	0 (0.0%)	1 (0.5%)

* Note: Discontinuation data is based on N=1 (0.8%) discontinuation within DTG/3TC treatment-naïve cohort and N=3 (1.6%) discontinuations within DTG/3TC SS cohort.

Conclusions

- TANDEM provides real world evidence supporting DTG/3TC use in the US.
- Almost all PLWH who initiated DTG/3TC experienced sustained virological suppression, with few treatment discontinuations.
- Primary reasons for initiation of DTG/3TC were to avoid long term toxicities (both cohorts), simplification/ streamlining of treatment (for SS sub-cohort) and convenience (for treatment-naïve sub-cohort).
- TANDEM reflects results from phase 3 clinical trials demonstrating DTG/3TC is an effective and well tolerated regimen when used in real-world settings in both treatment-naïve and SS PLWH.

Acknowledgments: The authors would like to thank the study investigators across each of the 24 US-based sites for their contribution towards the study in abstracting the data from medical charts of PLWH who were initiated on DTG/3TC.

References: 1. Waters, L. and Church, H. Curr. Opin. Infect. Dis. 2020; 33(1):28-33. 2. Cento, V. and Pento, C.F. JGAR. 2020; 20:228-237. 3. Katalam, C et al. AIDS. 2017; 31:1065-1071. 4. Cahn, P. et al. Lancet. 2019; 393(10167):143-155

This content was acquired following an unsolicited medical information enquiry by a healthcare professional. Always consult the product information for your country, before prescribing a ViiV medicine. ViiV does not recommend the use of our medicines outside the terms of their licence. In some cases, the scientific Information requested and downloaded may relate to the use of our medicine(s) outside of their license.