P-1021

The Evolution of Integrase Strand Transfer Inhibitors: VH-184, a Third-Generation Inhibitor With an Enhanced Resistance Profile

Jerry L. Jeffrey,[1] Jon Collins,[1] Luise Rogg,[2] Joseph Horton,[3] Sherene Min,[1] Cynthia Donovan,[1] Turner Overton,[1] Jorge Villacian[4]

[1]ViiV Healthcare, Durham, NC, USA; [2]GSK, Durham, NC, USA; [3]Parexel International, Durham, NC, USA; [4]ViiV Healthcare, London, UK

Presenting author:
Turner Overton
410 Blackwell Street
Durham, NC 27701
edgar.t.overton@viivhealthcare.com
205-842-4021

Please scan the QR code for additional resources

Key Takeaways

- **VH4524184 (VH-184) is a novel third-generation integrase strand transfer inhibitor (INSTI) with long-acting potential that has demonstrated highly potent antiviral activity similar to dolutegravir (DTG)**
- **In vitro data demonstrate the enhanced resistance profile of VH-184, showing activity against first- and second-generation INSTI-resistant strains**
- **VH-184 was developed to combine the potency of second-generation INSTIs with an enhanced resistance profile and long-acting potential to support durable treatment over time**

Plain Language Summary

- INSTIs are an important type of medicine for the treatment of HIV
- VH-184 is an experimental INSTI being developed by ViiV Healthcare; in laboratory studies, VH-184 was active against HIV that had become resistant to other INSTIs
- In a study where VH-184 was taken by mouth, it reduced the amount of HIV in the blood and was well tolerated
- VH-184 can also stay in the body for a long time, which means it might only need to be given twice a year as an injection

Introduction

- INSTIs have transformed antiretroviral therapy (ART) by offering durable potency and favorable tolerability
- Although first-generation INSTIs raltegravir (RAL) and elvitegravir (EVG) were an important innovation when first introduced, they have a low-to-moderate barrier to resistance[1]
- In the modern treatment era, second-generation INSTIs, including DTG, bictegravir (BIC), and cabotegravir (CAB), offer a higher barrier to resistance than RAL or EVG
  - However, the widespread use of second-generation INSTIs increases selective pressure which, together with virus heterogeneity, could impact long-term durability and drive emergent resistance, underscoring the need for next-generation INSTIs with an even higher barrier to resistance
- VH-184 is a third-generation INSTI with an enhanced resistance profile, showing activity against first- and second-generation INSTI-resistant strains in vitro[2,3]
- In resistance selection studies, no resistance mutations were identified through 112 days of selective pressure with VH-184[3]
- VH-184 monotherapy demonstrated highly potent antiviral activity similar to DTG in a proof-of-concept study, with no treatment-emergent INSTI resistance detected[4]
- VH-184 has long-acting potential, with twice-yearly dosing achievable based on available data[5]
- Here, we present the evolution of the INSTI class, including findings characterizing the enhanced resistance profile of VH-184


INSTI Evolution: From DTG to VH-184

In Vitro Resistance Profile of DTG

Antiviral Activity of DTG Against Viruses With Signature First-Generation INSTI Mutations

- Resistance selection studies and experiments of the antiviral activity of DTG against first-generation INSTI-resistant molecular clones established the resistance profile of DTG[6]
- No highly resistant mutants were isolated through Day 112 in resistance selection studies with DTG; substitutions that emerged conferred a fold change (FC; maximum, 4.1-fold) in the half-maximal effective concentration (EC_{50})[6]
- DTG retained activity against all single INSTI-resistant mutants tested, including those in the Y143, N155, and Q148 pathways (Figure 1)[6]
  - RAL and EVG showed FCs of up to >1700-fold against these same variants

Figure 1. Mean FC Against Molecular Clones With Key INSTI Substitutions[6]



Note that some data extend beyond the maximum mean FC of 50. [a]Each value represents a mean of 3 to 5 independent experiments performed in duplicate. [b]Mean FC value from 1 or 2 experiments.

- Only 2 double mutants (E138K/Q148K and Q148R/N155H) reduced DTG susceptibility ≥10-fold; notably, the single mutations driving these pathways were not selected during DTG passage[6]
- These findings demonstrated that DTG retained potent activity against HIV-1 variants harboring first-generation INSTI-resistant mutations, with a resistance profile markedly distinct from that of RAL and EVG[6]


Findings from in vitro studies established DTG as a second-generation INSTI with potent activity against signature first-generation INSTI-resistant mutants

Phase 3 Trials of DTG: VIKING-3 and VIKING-4

Results From VIKING-3 and VIKING-4 Confirmed Clinical Activity of DTG in Adults Who Were Heavily Treatment Experienced With INSTI-Resistant HIV-1

- The phase 3 VIKING-3 (single-arm, open-label; N=183) and VIKING-4 (randomized, double-blind, placebo-controlled; N=30) studies evaluated DTG 50 mg twice daily in adults with multi-class drug resistance, including resistance to RAL and/or EVG, on a failing regimen[7,8]
- In VIKING-3, 63% of participants achieved HIV-1 RNA <50 c/mL at Week 48 when receiving DTG twice daily with optimized background regimen despite extensive resistance[9]
- VIKING-4 confirmed the antiviral activity of DTG, with DTG functional monotherapy demonstrating a significant reduction in HIV-1 RNA vs placebo at Day 8 (treatment difference, −1.16 $\log_{10}$ c/mL; P<0.001)[8]
- Virologic response in both VIKING-3 and VIKING-4 was associated with baseline integrase genotype and was most attenuated in participants harboring Q148 mutations with accumulating secondary substitutions (Figure 2)[8,9]

Figure 2. Proportions of Participants in VIKING-3 and VIKING-4 Achieving Virologic Suppression (Snapshot HIV-1 RNA <50 c/mL) on DTG 50 mg Twice Daily + OBR at Week 48[8,9]



OBR, optimized background regimen; RAM, resistance-associated mutation. [a]Among other baseline virus mutation subgroups, HIV-1 RNA <50 c/mL at Week 48 was 100% (1/1) in the T66 subgroup, 79% (22/28) in the Y143 subgroup, 82% (27/33) in the N155 subgroup, and 50% (4/8) in the subgroup with ≥2 primary mutations. [b]Including L74I, E138A/K/T, and G140A/C/S.


VIKING-3 and VIKING-4 established the high barrier to resistance of DTG, demonstrating efficacy in participants with first-generation INSTI resistance

In Vitro Resistance Profile of VH-184

Findings Support VH-184 as a Third-Generation INSTI With an Enhanced Resistance Profile

- In an early in vitro experiment, VH-184 retained activity against all single INSTI-resistant mutants tested, including those in the Y143, G118, N155, and Q148 pathways, as well as the R263K mutation[3]
- In another study,[10] the resistance profile of VH-184 was evaluated against a panel of second-generation INSTI-resistant HIV-1 pseudotyped virus populations and clonal variants derived from participants meeting confirmed virologic withdrawal criteria in the phase 3 SAILING[11] (n=7; Week 132) and phase 3b DAWNING[12] (n=7; Week 216) studies
  - The proportion of virus populations and clonal variants achieving inhibitory quotients for VH-184 coverage against clonal strains was evaluated by combining exposure, viral potency, and clonal FC to predict efficacy of oral VH-184 45 mg
- VH-184 showed robust coverage across variant genotypes with key second-generation INSTI resistance mutations (Figure 3)

Figure 3. Heat Map Depicting the Proportion of Individuals Achieving Each IQ With VH-184 for Each Respective Clonal Variant Strain[10]




C_{trough}, plasma drug concentration at the end of the dosing interval; IQ, inhibitory quotient; PA-EC_{90}, protein-adjusted 90% effective concentration. [a]The IQ represents the ratio of drug exposure to viral susceptibility; IQ = (plasma drug concentration, C_{trough})/(PA-EC_{90} × ClonalFC). C_{trough} based on oral VH-184 45 mg.

VH-184 showed an enhanced resistance profile, retaining activity against INSTI-resistant pseudotyped viruses

INSTI Evolution and VH-184 Development

- The INSTI class has evolved from the first-generation INSTIs RAL and EVG to the second-generation INSTIs DTG, BIC, and long-acting CAB; VH-184 is the first third-generation INSTI to enter clinical development
- VH-184 retains potent antiviral activity against a broad panel of first- and second-generation INSTI-resistant variants
- Oral VH-184 monotherapy demonstrated highly potent antiviral activity in a proof-of-concept study, with up to a mean reduction in viral load of 2.31 $\log_{10}$ c/mL and no resistance to VH-184 detected[4]
- VH-184 represents the next step in the evolution of the INSTI class (Figure 4), showing
  - An enhanced resistance profile[2]
  - Potency similar to DTG with no treatment-emergent INSTI resistance during monotherapy[4]
  - Potential for extended dosing intervals[5]

Figure 4. Summary of the Evolution of the INSTI Class, Leading to the Third-Generation INSTI VH-184



RAM, resistance-associated mutation.

Conclusions

- Together, these data further support the development of VH-184 as a core agent in a complete long-acting regimen
- VH-184 will be further evaluated among people naive to ART in the phase 2b INNOVATE study (NCT07202546); an upcoming study in people with INSTI resistance is also planned

Acknowledgments: These studies were funded by ViiV Healthcare. Editorial assistance and graphic design support for this poster were provided under the direction of the authors by Fingerpaint Medical and funded by ViiV Healthcare.

References: 1. Doyle et al. *J Antimicrob Chemother.* 2015;70:3080-3086. **2.** Rogg et al. *Clin Infect Dis.* 2025;81:510-520. **3.** Seki et al. AIDS 2024; Munich, Germany. Poster WEPEB114. **4.** Rogg et al. *Clin Infect Dis.* 2026 [Epub ahead of print]. **5.** Back et al. CROI 2026; Denver, CO. Oral presentation 176. **6.** Kobayashi et al. *Antimicrob Agents Chemother.* 2011;55:813-821. **7.** Castagna et al. *J Infect Dis.* 2014;210:354-362. **8.** Akil et al. *Antivir Ther.* 2015;20:343-348. **9.** Vavro et al. European Meeting on HIV & Hepatitis 2014; Barcelona, Spain. Abstract O-10. **10.** Underwood et al. CROI 2026; Denver, CO. Poster 554. **11.** Cahn et al. *Lancet.* 2013;382:700-708. **12.** Aboud et al. *Lancet Infect Dis.* 2019;19:253-264.